Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

20862

[redacted] 50X1

DATE: January 30, 1953

SUBJECTS: New Products - January, 1953

[redacted] 50X1

Biomydrin

Description:	Aqueous solution with a pH of 6.2 containing:	
	neomycin sulfate	0.1%
	gramicidin	0.005%
	thonzylamine hydrochloride	1.0%
	phenylephrine hydrochloride	0.25%
	thonzonium bromide	0.05%
Indications:	For therapy of a wide range of common upper respiratory disorders and allergic conditions including rhinitis, sinusitis, nasopharyngitis as well as acute bacterial infections of the nasal mucosa.	
Administration:	Adult: 2 or 3 sprays in each nostril, four or five times a day as directed by physician. Children: 1 or 2 sprays in each nostril, four or five times a day as directed by physician.	

50X1 [redacted]
50X1
50X1 [redacted]

Gantricillin

50X1 Description:	Each tablet contains:	
50X1	Gantrisin (3,4-dimethyl-5-sulfanilamido-isoxazole)	5 mg.
	Crystalline penicillin G potassium	100,000 u.

50X1 [redacted]
50X1

Myciguent Cream

Description:	Each Gram contains:	
	neomycin sulfate	5 mg.
Indications:	Skin infections as impetigo, barber's itch, infectious exzematoid dermatitis, pustules, pustular acne, infected ulcers and boils.	
Administration:	Two to five times daily	

50X1 [redacted]

Neosone Ophthalmic Ointment

Description:	Each gram contains:	
	Cortisone Acetate	15 mg.
	Neomycin Sulfate	5 mg.

50X1 [redacted]

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Syncrobin Suppositories

Description:	Each suppository contains:	
	Penicillin G Potassium	200,000 u.
	Dihydrostreptomycin	200 mg.
Indications:	For treatment of a wide range of pathogenic disorders of the vaginal tract and for preoperative and postoperative use in major gynecologic surgery, and for controlling penicillin-sensitive and streptomycin-sensitive secondary invaders often associated with trichomonas vaginitis eukorrhea, and other disorders of the vaginal tract.	

50X1

Tyotocin Ear Drops

Description:	Formula for Tyotocin is:	
	Tyrothricin	0.05%
	benzocaine	1.25%
	antipyrine	5.0%
	hexylresoricinol	0.1%
	anhydrous propylene glycol	50.0%
	anhydrous glycerin q.s.	100.0%
	alcohol	0.5%
Indications:	In the treatment of mycotic and bacterial infections of the external auditory canal, such as otitis externa, furunculosis and secondarily infected exzema. Also for relief from pain and congestion in acute otitis media when the ear drum is intact.	
Administration:	May be instilled directly into the alcohol-cleansed aural canal with a dropper, or if preferred, the aural canal may be packed with guaze wick saturated with Tyotocin. Wick should be kept moist and removed 12 to 24 hours later.	

50X1

Tyrophyll Troches

Description:	Each troche contains:	
	tyrothricin	2 mg.
	benzocaine	7.5 mg.
	chlorophyllin	2 mg.
Indications:	For the relief and treatment of throat infections and irritations.	
Administration:	1 troche every two hours - not more than 8 troches in a 24-hour period.	

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Foreign

Mylipen

Description: Each mil contains 300,000 units procaine penicillin G suspended in isotonic solution; the suspension also contains about 1.5% dispersing agents and 0.025% cetrimide as bacteriostat.

Indications: Infections due to penicillin-sensitive organisms when a prolonged inhibitory level in the blood is desired.

Administration: Intramuscularly 1 mil every 24 hours

50X1

Mylipen Cream

Description: Each gram contains:
Procaine Penicillin 10,000 u.

Indications: Superficial skin diseases caused by penicillin-sensitive organisms; pyogenic dermatoses, ecthyma, impetigo, sycosis barbae, otitis externa, secondarily infected eczema and other conditions secondarily infected with staphylococci prophylaxis and treatment of infection in burns.

Administration: Apply to affected area three or four times daily and cover if necessary.

50X1

Neobiocin

Description: Neomycin sulfate
Sodium Penicillin G
Dihydrostreptomycin

Indications: Gram positive and gram negative bacteria

Administration: Intramuscular. Will maintain appreciable blood level for 24 hours - injections cannot be prolonged for more than six days.

50X1

Penicillin Monad Tulle

Description: Sterilized gauze dressing of wide mesh is impregnated with an emulsifying base containing 1,000 units of penicillin per gram, equivalent to 160 units of penicillin per square inch of tulle.

Indications: an effective bacteriostatic dressing for operation wounds, including those of eyes, ears, and nose, and those of skin grafting.

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Penicillin for Eye Drops

50X1 [redacted] introduction [redacted] strength of 15,000 i.u. per
50X1 product of Tabloid hypodermic benzylpenicillin
50X1 with sodium citrate. This product is in-
50X1 tended for extemporaneous preparation of the
50X1 penicillin eyedrops of the B.P.C. and the N.F.
50X1 and replaces Tabloid penicillin hypodermic of
50X1 the same strength. [redacted]

Spemycine

Description: Each dose contains

Dihydrostreptomycin	.5 gm.
Sodium Penicillin G	200,000 U.
Procaine Penicillin G	300,000 U.

Indications: Severe throat infections, phlegmons, infected wounds and suppurating lesions, staphylococcal infections, streptococcal infections, pneumococcal infections, gonococcal infections, and their complications pneumococcal infections, broncho-pneumonia, purulent pleurisy, respiratory complications of infectious diseases, otitis and mastoiditis, infections of the pulmonary tract, acute peritonitis (in connection with surgical treatment), pre-and-post operative treatment for intestinal surgery, urinary infections.

Administration: 1 injection every 24 hours.

50X1 [redacted]

Trulfacillin

3-200 Suspension

Description: Each 5cc. teaspoonful contains:

Sulfamethazine	65 mg.
Sulfadiazine	65 mg.
Sulfamerazine	65 mg.
(A total of 3 gr. of sulfonamide)	
Benzethacillin	200,000 Int. U.
(Dibenzylethylenediamine dipenicillin-G)	

Administration: Infants and Children: 1 teaspoonful per 4 pounds of body weight per day, in divided doses.

Form Supplied: Bottles of 60 cc.

7½-100 Suspension

Description: Each 5 cc. teaspoonful contains:

Sulfamethazine	.167 G.
Sulfadiazine	.167 G.
Sulfamerazine	.167 G.
(A total of 7½ gr. of sulfonamide)	
Benzethacillin	100,000 Int. U.
(Dibenzylethylenediamine dipenicillin-G)	

Administratio
Administration: 1 to 2 teaspoonfuls every 6 to 8 hours.

50X1 [redacted]

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

7½-100 Tablets

Description? Each tablet contains:

Sulfamethazine	.167 G
Sulfadiazine	.167 G
Sulfamerazine	.167 G
(a total of 7½ gr. of sulfonamides)	
Potassium Penicillin G crystalline	100,000 U.I.

50X1 Administration: 1 to 2 tablets every 4 to 8 hours

50X1

3-50 Tablets

Description: Each tablet contains:

Sulfamethazine	64 mg.
Sulfadiazine	64 mg.
Sulfamerazine	64 mg.
(a total of 3 gr. of sulfonamides)	
Potassium Penicillin G Crystalline	50,000 U.I.

Administration: 1 tablet per day per 4 pounds of body weight in divided doses.

50X1

5-25 Tablets

Description: Each tablet contains:

Sulfamethazine	.1 G
Sulfadiazine	.1 G
Sulfamerazine	.1 G
(a total of 5 gr. of sulfonamides)	
Potassium Penicillin G Crystalline	25,000 U.I.

50X1 Administration: 1 to 4 tablets every 4 hours followed by 1 to 2 tablets every 4 hours

50X1

Indications: (for all forms of Trulfacillin) Used in treating pneumococcus infections, staphylococcus infections, hemolytic streptococcus infections, gonorrhea, tonsillitis, scarlet fever, otitis media and Vincent's angina.

50X1

50X1

Tyrosolven

Description: Each lozenge contains tyrothricin and benzocaine with a tyrothricin solvent in a pleasantly flavored base.

Indications: Treatment of mouth and throat infections and is stated to act only in the mouth and throat and is not absorbed into the system.

50X1

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4

Tyrosolven (cont'd.)

50X1

Tyrosolvin - as a topical solution for treatment in opthamology, otolaryngology and dermatology. Supplied in 10mg. and 30mg. bottles with eyedropper.

50X1

Tyrosolvin - as an opthalmic cream with applicator in tubes of 10 gm. and 40 gm.

Tyrosolvetter - a Sugar tablet containing 1 mg. of tyrothrycin and 5 mg. of benzocaine. Supplied in packs of 25 and 50 for treatment of gingivitis, stomatitis, ulcers and prophylaxis.

50X1

Declassified in Part - Sanitized Copy Approved for Release @ 50-Yr 2013/07/23 : CIA-RDP83-00423R000700410005-4